The Turán density of the tight five-cycle

P. Graham
https://orcid.org/0009-0007-8493-1505

October 27, 2026

Abstract

The tight five-cycle is the three-uniform hypergraph whose edges are the five cyclically consecutive triples on five distinct vertices. We prove that its Turán density is $2\sqrt{3}-3$. This answers the conjecture usually attributed to Mubayi and Rödl, explicitly stated by Mubayi, Pikhurko and Sudakov. Mubayi and Rödl's recursive construction gives the lower bound. The upper bound follows the reduction of Bodnár, León, Liu and Pikhurko, whose theorem concerns forbidding both the tight four-cycle and the tight five-cycle, or a tight cycle of length at least seven not divisible by three. We establish the two estimates needed for the five-cycle alone by exact rational certificates, using weighted link inequalities to include configurations containing a complete four-vertex hypergraph. We give the finite counting argument that turns the certificates into these estimates and then into the density bound. The whole proof, including this counting argument, is checked in Lean; its finite computations run as compiled code, so the check trusts Lean's compiler as well as its kernel.

This work has not yet been reviewed by experts. The proofs, the Lean formalization and the text were produced by AI models (from OpenAI and Anthropic), working under the author's direction.

1 Introduction

A three-uniform hypergraph, or 3-graph, on V is a subset of $\binom{V}{3}$. Put

$$C_5^3 = \{\{i, i+1, i+2\} : i \in \mathbb{Z}/5\mathbb{Z}\}.$$

A copy in H means an injective map $f : \mathbb{Z}/5\mathbb{Z} \to V$ with $\{f(i), f(i+1), f(i+2)\} \in H$ for every i; extra edges are allowed. Write $f(n) = \mathrm{ex}(n, C_5^3)$ for the largest number of edges in a C_5^3-free 3-graph on n vertices, and $\alpha = 2\sqrt{3}-3$.

Theorem 1. *The injective, non-induced tight five-cycle has Turán density*

$$\pi(C_5^3) = \lim_{n\to\infty} \frac{\mathrm{ex}(n, C_5^3)}{\binom{n}{3}} = 2\sqrt{3}-3.$$

Precisely, for every real $\varepsilon > 0$ there is an integer $N \geq 3$ such that for every integer $n \geq N$,

$$(\alpha-\varepsilon)\binom{n}{3} \leq f(n) \leq (\alpha+\varepsilon)\binom{n}{3}.$$

Mubayi–Rödl's Construction 1 gives the lower bound [2, Construction 1][1]; the equality is stated explicitly by Mubayi, Pikhurko and Sudakov [3, §2.4, Conjecture 7]. They also record Razborov's

[1] Mubayi and Rödl credit Y. Kohayakawa for discussions leading to this construction [2, §7].

earlier upper bound $\pi(C_5^3) < 0.4683$ [3, 19, §2.4]. The usual attribution is recorded in Kamčev–Letzter–Pokrovskiy [4, Conjecture 1.1]; their Theorem 1.3 gives the equality for all sufficiently large cycle lengths not divisible by three. Bodnár–León–Liu–Pikhurko (BLLP) prove the equality for $\{C_4^3, C_5^3\}$ and for C_ℓ^3 with $\ell \geq 7$ and $3 \nmid \ell$ [5, Theorem 1.1].

Falgas-Ravry and Vaughan establish the sharp bound for 3-graphs whose edges are the triples inducing a directed out-star in a digraph [7, §3.1, Proposition 25 and Theorem 27]. Huang gives an analytic proof of the directed-star result [8, §§2 and 4]. This restricted class does not include every C_5^3-free 3-graph.

The partition method and reduction are BLLP's [5, Propositions 3.2–3.3, Lemma 4.1 and the proof of Theorem 2.4]. Here their two estimates are established for C_5^3 alone, with the precise hypotheses below. The difficult step is the estimate for bad edges minus missing triples when K_4^3 is allowed: BLLP identify this obstacle in [5, §7]. The additional link inequalities below permit exact certificates over all C_5^3-free six-vertex configurations. The colored flag setup and degree and local-maximality rows follow BLLP's supplementary calculations [6, notebook, zero-based code cells 1 and 4]. That notebook uses Bodnár's SageMath framework, documented as *FlagAlgebraToolbox* [10, §§1.2 and 2.4]. Finite sampling makes the certificate-to-graph passage explicit. The proof depends on these computations and their Lean formalization; Section 5 states the formal conclusion and its trust boundary. Earlier computations for C_5^3 alone include Xizhi Liu's numerical flag-algebra optimization [12, edge-density objective and saved output]. The present argument supplies exact certificates and the finite estimates needed for the unrestricted upper bound.

Ferudun's October 16, 2026 preprint [13, Theorems 1–3 and §5] optimizes the classical recursive construction by a logarithmic-state algorithm and determines its quadratic term. The recursive construction family is shared. Ferudun explicitly distinguishes this construction-side result from an upper bound for arbitrary C_5^3-free hypergraphs. The Ulam.AI UnsolvedMath record AIM-COMBINATORICS-0153 reports an exact deficit identity, weighted stability and an optimized finite recurrence within the same recursive family [14]. That summary also leaves the unrestricted upper bound open; its underlying proof was not read or checked here.

For the related cycle with one edge deleted, Balogh and Luo [16] determine the density for all sufficiently long lengths not divisible by three. Bodnár, León, Liu and Pikhurko [15, Theorem 1.1] and, independently, Lidický, Mattes and Pfender [17, Abstract] determine density 1/4. That is a different forbidden hypergraph.

2 Two partition estimates

For $S \subseteq V(H)$ let $T = V(H) \setminus S$ and define

$$\mathrm{cut}_H(S) = |\{e \in H : |e \cap S| = 2\}|, \qquad \mu(H) = \max_{S \subseteq V(H)} \mathrm{cut}_H(S),$$

$$B_H(S) = \{e \in H : |e \cap S| \in \{1,3\}\}, \qquad M_H(S) = \{e \in \tbinom{V}{3} \setminus H : |e \cap S| = 2\}.$$

A cut is locally maximal if $\mathrm{cut}_H(S \triangle \{v\}) \leq \mathrm{cut}_H(S)$ for every $v \in V(H)$. Thus a maximum cut is locally maximal. Write $d_H(v)$ for vertex degree, $\beta_0 = 4641/10000$ and $\beta_1 = 464101615/10^9$. The following are the C_5^3-only counterparts of BLLP's Propositions 3.2 and 3.3; they use binomial normalization, an asymptotic cut threshold, and coefficient one on the missing triples.

Proposition 2. *For every real $\varepsilon > 0$ there exists an integer N such that, for every integer $n \geq N$ and every C_5^3-free 3-graph H on n vertices, $|H| \geq \beta_0 \binom{n}{3}$ implies $\mu(H) \geq (2/5 - \varepsilon)\binom{n}{3}$.*

Proposition 3. *For every real $\xi > 0$ there exist a real $\delta > 0$ and an integer N such that, for every integer $n \geq N$, every C_5^3-free 3-graph H on n vertices and every $S \subseteq V(H)$, the hypotheses*

$$S \text{ is locally maximal}, \qquad d_H(u) \leq d_H(v) + \delta n^2 \quad \text{for all } u, v \in V(H),$$
$$\operatorname{cut}_H(S) \geq (2/5 - \delta)\binom{n}{3}, \qquad |H| \geq (\beta_1 - \delta)\binom{n}{3}$$

imply $|B_H(S)| - |M_H(S)| \leq \xi n^3$.

These hypotheses are auxiliary estimates, not restrictions in Theorem 1. In particular, the same cut of an extremal graph will meet both estimates.

3 Certificates and finite sampling

The certificates use Razborov's flag algebra method [18, §2] in the finite-sampling form described by BLLP [5, §3]. For $n \geq 12$, sample an ordered injection (x, u, v, a, b, d) into $V(H)$. Write xuv for the triple $\{x, u, v\}$ in edge indicators. If a cut is present, color its vertices by $c(z) = \mathbf{1}[z \notin S]$. Let q_K be the probability of its induced six-vertex isomorphism class K. For any function R of the sample, let $\bar{R}(K)$ be its average over all 6! orders of K. Then

$$\mathbb{E}R = \sum_K q_K \bar{R}(K), \qquad q_K \geq 0, \qquad \sum_K q_K = 1. \tag{1}$$

Only C_5^3-free classes occur, including classes containing K_4^3.

The link inequalities

Put $L_x(a, b) = \mathbf{1}[\{x, a, b\} \in H]$ and let $K_x(a, b, c)$ indicate that $\{x, a, b, c\}$ spans K_4^3. Put

$$C_x(a, c, d) = L_x(a, c)L_x(a, d)(1 - L_x(c, d))(1 - \mathbf{1}[\{a, c, d\} \in H]).$$

Repeated vertices are permitted in the following sums; an edge always has three distinct vertices.

Lemma 4. *If H is C_5^3-free, $x, a \in V(H)$, and $w_z \geq 0$ for every $z \in V(H)$, then*

$$\sum_{c,d} K_x(a, c, d)w_c w_d \leq \sum_{c,d} C_x(a, c, d)w_c w_d. \tag{2}$$

Multiplication by w_a and summation over a gives the corresponding three-variable inequality. Dropping the last non-edge factor on the right gives a further valid inequality.

Proof. For fixed x, a, let $F(b, c) = K_x(a, b, c)$ and $s_c = \sum_b F(b, c)w_b$. If $F(b, c) = F(b, d) = 1$, both $\{x, c, d\}$ and $\{a, c, d\}$ are absent: otherwise respectively the orders (c, a, b, d, x) and (x, b, c, a, d) give a tight five-cycle. Consequently $s_b \leq m_c := \sum_d C_x(a, c, d)w_d$ whenever $F(b, c) = 1$. Symmetry and $s_b/s_c + s_c/s_b \geq 2$ give

$$\sum_{b,c} F(b, c)w_b w_c \leq \sum_{b,c} F(b, c)w_b w_c \frac{s_b}{s_c} \leq \sum_c w_c m_c.$$

Terms of zero weight contribute zero; all denominators in a nonzero term are positive. This proves (2). □

Fixing roots, restrict each free vertex by the same predicate on its edges to the roots and its color. Applying Lemma 4 with indicator weights gives the three-variable rows, with or without the last non-edge factor, and the two-variable rows. Nonnegative rational weights give the weighted two-variable rows. On passing to distinct samples the diagonal is removed, producing an error at most $1/(n-5)$, or $W^2/(n-5)$ for two-variable weights bounded by W. The diagonal is retained before this passage; it need not vanish on K_4^3. Appendix A gives the other finite errors.

The cut certificate

A selector has a specified edge or non-edge type on (x, u, v) and a rational function p assigning each possible three-edge pattern of an external vertex a probability in $[0, 1]$. Given the roots, put each other vertex in a set U independently with that probability. Let D_s be the probability of the root type, and let N_s be the probability of that type together with $\{a, b, d\} \in H$ and $|\{a, b, d\} \cap U| = 2$. On a class, $\bar{N}_s(K)$ and $\bar{D}_s(K)$ denote the averages of these same six-vertex integrands. Averaging the number of selected edges gives

$$N_s \le \frac{6 D_s \mu(H)}{(n-3)_3}. \tag{3}$$

Indeed, for every fixed root triple and every outcome U, the selected edges avoiding the roots number at most $\mathrm{cut}_H(U) \le \mu(H)$; each edge has six orders.

The rational certificate verifies, on every uncolored class,

$$\sum_s \lambda_s (\bar{N}_s - \tfrac{2}{5}\bar{D}_s) = 10^{-4} + \sum_i y_i \bar{R}_i + \sum_j \langle Q_j, A_j \rangle + h_K. \tag{4}$$

Here $\lambda_s, y_i, h_K \ge 0$, $\sum_s \lambda_s = 1$, and every Q_j is positive semidefinite. The matrices A_j are products of rooted flag indicators; $\langle Q, A \rangle = \sum_{r,t} Q_{rt} A_{rt}$. The scalar rows are the link rows above and the two density products $(\mathbf{1}[xuv \in H] - \beta_0)\mathbf{1}[abd \in H]$ and its non-edge counterpart. All coefficients are averaged as in (1).

Proof of Proposition 2. Average (4) using (1). The finite errors in Appendix A, weighted by the fixed certificate coefficients, have total at most $C/(n-5)$ for all sufficiently large n, where C depends only on the certificate. Thus some selector satisfies $N_s - \frac{2}{5}D_s \ge 10^{-4} - C/(n-5) > 0$ for all sufficiently large n. Since $0 \le N_s \le D_s$, its denominator is positive. Equation (3) now gives $\mu(H) \ge \frac{2}{5}\binom{n-3}{3}$. Finally $\binom{n-3}{3}/\binom{n}{3} \to 1$, proving the stated quantifiers. □

The bad-minus-missing certificate

For a colored class set $t_K = (|B_K| - |M_K|)/20$. The second rational certificate verifies, class by class,

$$t_K + \sum_j \langle Q_j, M_j(K) \rangle + \sum_i y_i \bar{a}_i(K) + \sum_j e_j \bar{r}_j(K) \le 0. \tag{5}$$

Here Q_j are positive semidefinite, $y_i \ge 0$, and e_j may have either sign. The rows a_i express local maximality, cut density, edge density at β_1, and the three-variable link inequality without the last non-edge factor. The rows r_j express degree differences. Their explicit sampling functions are in Appendix A. Only the matrices and multipliers are divided by the certificate's positive integer scale; t_K is unscaled.

Proof of Proposition 3. Each triple occurs equally often in six-sets, so $\sum_K q_K t_K = (|B_H(S)| - |M_H(S)|)/\binom{n}{3}$ exactly. For hypotheses with slack δ', averaging (5) and using the finite errors gives

$$\frac{|B_H(S)| - |M_H(S)|}{\binom{n}{3}} \le A_n\delta' + K_n, \qquad A_n \longrightarrow A \in [0,\infty), \quad K_n \longrightarrow 0.$$

The absolute values of e_j are used for degree-difference errors. The density and cut products have lower bounds $-\rho_n\delta' - \eta_n$, with $\rho_n \to 1$, $\eta_n \to 0$ as defined in the appendix. All other errors are $O(1/n)$. Given $\xi > 0$, take $\delta = \min\{1, \xi/(A+1)\}$ and then N so that $A_n \le A+1$ and $K_n \le \xi$ for every $n \ge N$. The result follows from $2\xi\binom{n}{3} \le \xi n^3$. □

4 Proof of the density theorem

Proof of Theorem 1. For the lower bound use Mubayi–Rödl's construction [2, Construction 1]: partition into A, B, take every triple with two vertices in A and one in B, and repeat inside B (Figure 1). A pair shared by two edges determines their construction level, so all edges of a tight cycle would have the same level. Their three consecutive vertices would then have a periodic part pattern with two A's and one B, incompatible with length five [4, Example 1.2]. With $|B|/n \to r = (\sqrt{3}-1)/2$, the limiting density is $3(1-r)^2 r \sum_{j\ge 0} r^{3j} = \alpha$; rounding and a finite-depth approximation give $f(n) \ge (\alpha - o(1))\binom{n}{3}$.




Figure 1: The recursive construction. The triangle illustrates an edge with two vertices in A and one in B; all such triples are included.

For the upper bound let H attain $f(n)$. Replacing a vertex u by a clone of v preserves C_5^3-freeness: the collapsing map sends edges to edges, and every pair of vertices of C_5^3 lies in an edge, so any cycle image is still injective. Extremality therefore gives $d_H(v) \le d_H(u) + n$ for all u, v; the possible loss is at most the codegree of u, v. Choose a maximum cut S, which is locally maximal, and set $s = |S|$, $m = n - s$. Since $\alpha > \beta_0$ and $\alpha \ge \beta_1$, the lower bound, Proposition 2, and the degree inequality meet all the hypotheses of Proposition 3 on this same cut for sufficiently large n. For each fixed $\xi > 0$ we obtain

$$f(n) \le \binom{s}{2} m + f(m) + \xi n^3 \le \tfrac{1}{2} s^2 m + f(m) + \xi n^3. \tag{6}$$

Indeed, the edges consist of the cut edges, the bad edges and the edges inside $V(H) \setminus S$; the cut edges number $\binom{s}{2}m - |M_H(S)|$. We may use Proposition 2 with error at most $1/5$. For large n, $\binom{n}{3} \ge n^3/12$, so $\mathrm{cut}_H(S) \ge n^3/60$. On the other hand $\mathrm{cut}_H(S) \le s^2 n/2$, which rules out $s < n/6$. Hence $m \le 5n/6$.

The scalar inequality in BLLP [5, Fact 2.9(i)] is

$$\frac{s^2 m}{2} \le \frac{\alpha}{6}\left((s+m)^3 - m^3\right) \quad (s, m \ge 0). \tag{7}$$

It follows from the factorization

$$\frac{\alpha}{6}\left((s+m)^3 - m^3\right) - \frac{s^2 m}{2} = \frac{\alpha s}{6}(s - \sqrt{3}m)^2 \ge 0.$$

Let $D = \limsup_{n\to\infty} f(n)/\binom{n}{3} \in [0,1]$. For every $\eta > 0$ there is a constant K_η such that $f(m) \le (D+\eta)m^3/6 + K_\eta$ for every integer $m \ge 0$. If $D > \alpha$, combine (6)–(7) and $m^3 \le (5/6)^3 n^3$ to obtain

$$\frac{6f(n)}{n^3} \le \alpha + (D+\eta-\alpha)(5/6)^3 + 6\xi + \frac{6K_\eta}{n^3}$$

for all sufficiently large n. Taking a subsequence tending to the limsup and then letting η, ξ tend to zero gives $D - \alpha \le (5/6)^3(D-\alpha)$, a contradiction. Together with the lower bound this proves convergence and the theorem. □

5 Formal verification

The closed Lean theorem has the following name and type, with $\mathbb{N}$ and $\mathbb{R}$ denoting Lean's natural and real number types:

```
TightCycleC5.Bridge.mubayi_rodl_checked :
Filter.Tendsto
  (fun n :  ℕ =>
    (TightCycleC5.ex n :  ℝ) / (n.choose 3 :  ℝ))
  Filter.atTop (nhds (2 * Real.sqrt 3 - 3))
```

Here `TightCycleC5.ex n` is the maximum edge count over all finite sets H of three-element subsets of `Fin n` for which there is no injection f : `Fin 5` $\hookrightarrow$ `Fin n` with $\{f(i), f(i+1), f(i+2)\} \in H$ for every i; addition in `Fin 5` is cyclic. Thus it is exactly $f(n)$ from Theorem 1, with no induced-copy restriction or assumption on H or its partitions. Total real division defines the initial terms for $n < 3$, which do not affect the limit.

The class-by-class certificate assertions (4) and (5), including positivity and class coverage, are checked in the formalization and by the accompanying exact certificate checkers; they are not derived from a printed list of coefficients here. The written argument proves their passage to Propositions 2–3 and Theorem 1. The formal proof includes this passage and the density reduction. It depends on 692 `native_decide` facts, trusting Lean's compiler and runtime as well as its kernel; it is not a kernel-only proof. These facts check finite arithmetic, table and coefficient comparisons, and two rational Gram identities. The class covers, arithmetic-range coverage, all six uncolored positive-semidefinite matrices and the other 36 colored matrices have kernel proofs.

The accompanying Lean bundle is unreleased at the time of writing. Its checking material [1] specifies the pinned toolchain, all standard axioms, the exact 692-fact dependency list and reproduction commands. The recorded full replay checked the frozen mathematical sources; a later adapter check reused those validated dependencies and elaborated a fresh probe of the theorem's exact type and allowed axioms. That probe is not a second replay. The original endpoint placeholder supplies only a type: its proof is outside the checked theorem's dependency chain. The checking material distinguishes these executions from the standalone Lake build, which was not exercised.

For related formalization, Jeong, Park, Hyun, Oum and Yang [9, Abstract] formalize flag algebras for finite simple graphs. Jeong, Park, Im, Lee and Yang combine degree-stationarity with an exact seven-vertex certificate for the tetrahedron and formalize its bound [11, §§2.5 and 3]. The present development proves the finite sampling estimates for the 3-graphs and colored 3-graphs used here directly.

Appendix A Finite sampling estimates

Here are the row definitions and estimates used in the two certificate arguments. They apply for $n \geq 12$ and always to ordered injections; $(n)_k = n(n-1)\cdots(n-k+1)$. A flag is the induced graph, its colors when present, and its designated ordered roots, with free vertices identified up to the specified permutations. Its indicator vector is one-hot; a coarsening of such a flag is also permitted.

Link rows. For roots (x, u, v), let $p(z) = (\mathbf{1}[xuz \in H], \mathbf{1}[xvz \in H], \mathbf{1}[uvz \in H])$. Choose a nonempty subset of its eight patterns, and optionally a set of allowed colors. Every one of a, b, d satisfies the same restriction; a root-type indicator may also be imposed. The row is the right side minus the left side of the three-variable version of Lemma 4, with or without the last non-edge factor. Its average is at least $-1/(n-5)$: the removed diagonal has at most $(n-3)(n-4)$ terms and the conditional denominator is $(n-3)_3$. For the two-variable rows fix (x, u, v, a) and use a common nonnegative weight on each free vertex determined by

$$(\mathbf{1}[xuz \in H], \mathbf{1}[xvz \in H], \mathbf{1}[uvz \in H], \mathbf{1}[auz \in H], \mathbf{1}[avz \in H]).$$

If its maximum is W, the diagonal weighs at most $(n-4)W^2$ and the average is at least $-W^2/(n-5)$.

Positive semidefinite rows. For a fixed root tuple θ of a specified type, let f_θ be a one-hot flag vector and Q be positive semidefinite. Put $m_Q = \max_i Q_{ii}$. Summing all pairs of flags gives $X^T Q X \geq 0$, where X is the sum of their indicator vectors. Removing intersecting free sets gives respectively the lower bounds

$$-\frac{m_Q}{n-5} \quad \text{(four roots)}, \qquad -\frac{(4n-14)m_Q}{(n-4)(n-5)} \quad \text{(two roots)}.$$

For two roots there are $(n-2)(n-3)(4n-14)$ intersecting pairs of ordered free pairs, and $|Q_{ij}| \leq \sqrt{Q_{ii}Q_{jj}} \leq m_Q$. For four roots only the $n-4$ diagonal pairs are removed.

Products on disjoint triples. For functions g, h of ordered triples, $|g|, |h| \leq 1$, let $\mathbb{E}_3$ denote their average on a uniformly sampled ordered triple of distinct vertices. Then

$$|\mathbb{E}[g(x,u,v)h(a,b,d)] - \rho_n \mathbb{E}_3 g\, \mathbb{E}_3 h| \leq \eta_n, \qquad \rho_n = \frac{(n)_3}{(n-3)_3}, \quad \eta_n = \frac{9n^2 - 45n + 60}{(n-3)_3}.$$

This follows by removing the pairs of triples that intersect, since $(n)_3 - (n-3)_3 = 9n^2 - 45n + 60$. For cut rows use $g = \mathbf{1}[xuv \in H]\mathbf{1}[c(x)+c(u)+c(v) = 1] - 2/5$; for density rows use $g = \mathbf{1}[xuv \in H] - \beta_i$. The multiplier h is a three-vertex flag indicator (the edge and non-edge indicators in the uncolored certificate). Thus $\mathbb{E}_3 g \geq -\delta'$ gives the lower bound $-\rho_n\delta' - \eta_n$ when $h \geq 0$. The unconditional density row has lower bound $-\delta'$ exactly. In particular, $(n-5)\eta_n \to 9$; a uniform bound uses a constant strictly larger than nine.

Local maximality and regularity. Put

$$\ell(x,u,v) = \mathbf{1}[xuv \in H](\mathbf{1}[c(u)+c(v) = 1] - \mathbf{1}[c(u)+c(v) = 0])(1 - 2c(x)).$$

For every one-root flag indicator $I_F(x, a, b, d)$, local maximality gives

$$\mathbb{E}[\ell(x,u,v)I_F(x,a,b,d)] \geq -\frac{6(n-3)}{(n-4)(n-5)}.$$

For fixed x, the unrestricted sum of ℓ over ordered (u,v) is $2(\mathrm{cut}_H(S)-\mathrm{cut}_H(S\triangle\{x\}))\geq 0$. The sum of I_F over the free triple is nonnegative and independent of (u,v). Deleting triples meeting $\{u,v\}$ removes $6(n-3)^2$ possibilities per fixed pair, giving the displayed bound. For each two-root flag indicator $I_F(x,u,b,d)$, ordered degree regularity gives

$$|\mathbb{E}[(\mathbf{1}[xva\in H]-\mathbf{1}[uva\in H])I_F(x,u,b,d)]|\leq \frac{2\delta' n^2+4n-14}{(n-4)(n-5)}.$$

Indeed, summing over (v,a) outside $\{x,u\}$ gives $2(d_H(x)-d_H(u))$, because codegree terms cancel; at most $4n-14$ pairs meet $\{b,d\}$. Regularity is required for all ordered x,u, including equality, so both signs have this bound.

Certificate data. The first certificate has nine selectors and six positive semidefinite blocks: five with four roots and one with two. It uses the three-variable link rows with and without the non-edge factor, the two-variable indicator and rational-weight rows, and the two density products above. The second has 38 blocks: three with two roots and 35 with four. Its scalar rows comprise 80 local, eight cut, nine density and 12,750 three-variable link rows, and 120 signed regularity rows. Each matrix entry is the joint indicator of its two flags on the appropriate disjoint free sets; the matrix pairing includes both ordered off-diagonal terms. In every case coefficients are rebuilt by averaging all 720 orders of a class. With color 1 assigned to $V(H)\setminus S$, bad edges have color sum zero or two and missing triples have color sum one. This explains the exact target normalization by $\binom{6}{3}=20$. These specifications, together with the rational certificate entries, determine (4) and (5) without any numerical optimization premise.

References

[1] P. Graham, *Checking material for The Turán density of the tight five-cycle*, version 1, accompanying Lean bundle (unreleased). The formalization guide and certificate checkers accompany the paper; formalization guide. Toolchain: `leanprover/lean4:v4.32.2`; Mathlib revision 905b95818eb32af7874a58b427f50c1711a5e96c.

[2] D. Mubayi and V. Rödl, *On the Turán Number of Triple Systems*, Journal of Combinatorial Theory, Series A **100** (2002), no. 1, 136–152. Author's copy, Construction 1, p. 13: https://homepages.math.uic.edu/~mubayi/papers/hypturan.pdf.

[3] D. Mubayi, O. Pikhurko and B. Sudakov, *Hypergraph Turán Problem: Some Open Questions*, October 10, 2011, §2.4, Conjecture 7, p. 3. https://rose-kiwi.lnx.warwick.ac.uk/E/MubayiPikhurkoSudakov11aim.pdf.

[4] N. Kamčev, S. Letzter and A. Pokrovskiy, *The Turán Density of Tight Cycles in Three-Uniform Hypergraphs*, International Mathematics Research Notices **2024** (2024), no. 6, 4804–4841. Numbering follows arXiv:2209.08134v3 (October 6, 2023; first posted September 16, 2022). https://arxiv.org/abs/2209.08134v3.

[5] L. Bodnár, J. León, X. Liu and O. Pikhurko, *The Turán density of short tight cycles*, arXiv:2506.03223v1 (first posted June 3, 2025). All numbered references follow the versioned PDF. https://arxiv.org/abs/2506.03223v1.

[6] L. Bodnár, J. León, X. Liu and O. Pikhurko, *The Turán number of* $\{C_5,K_4\}$, supplementary notebook, introduction and zero-based code cells 1–6, June 3, 2025. https://github.com/bodnalev/supplementary_files/blob/5766324d010da24a66271c0fbe38460cd377985b/no_cl/no_c5_k4.ipynb.

[7] V. Falgas-Ravry and E. R. Vaughan, *Turán H-densities for 3-graphs*, Electronic Journal of Combinatorics **19** (2012), no. 3, P40. Numbering follows arXiv:1201.4326v1 (first posted January 20, 2012). https://arxiv.org/abs/1201.4326v1.

[8] H. Huang, *On the maximum induced density of directed stars and related problems*, arXiv:1303.3101v2 (March 17, 2013; first posted March 13, 2013). https://arxiv.org/abs/1303.3101v2.

[9] G. Jeong, S. Park, J. Hyun, S.-i. Oum and H. Yang, *Formalizing Flag Algebras in Lean*, arXiv:2607.23500v2 (October 14, 2026; first posted July 26, 2026). https://arxiv.org/abs/2607.23500v2.

[10] L. Bodnár, *FlagAlgebraToolbox: Flag Algebra Computations in SageMath*, arXiv:2601.06590v3 (September 12, 2026; first posted January 10, 2026). This version documents release 0.3; the earlier supplementary notebook [6] does not identify a release tag. https://arxiv.org/abs/2601.06590v3.

[11] G. Jeong, S. Park, S. Im, J. Lee and H. Yang, *A New Upper Bound for the Turán Density of the Tetrahedron*, arXiv:2609.27495v2 (October 2, 2026; first posted September 23, 2026). https://arxiv.org/abs/2609.27495v2.

[12] X. Liu, *C53_count.ipynb*, edge-density objective and saved output, February 6, 2026, repository revision `1b5c8b15c545ea15587f8240bd2149a29aaba857`. https://github.com/xliu2022/xliu2022.github.io/blob/1b5c8b15c545ea15587f8240bd2149a29aaba857/FlagAlgebra_Certificate/HypergraphGeneralizeTuran/script/C53_count.ipynb.

[13] A. Ferudun, *An Exact Logarithmic-State Algorithm for Recursive Tight-Cycle Constructions*, version 1.0, October 16, 2026. https://doi.org/10.5281/zenodo.23421447.

[14] Ulam.AI, *UnsolvedMath*, record AIM-COMBINATORICS-0153 (record id 20001028), research summary on recursive tight-cycle constructions. The record metadata is dated August 14, 2026. Only the pinned dataset record was read, not the underlying attempt or proof. https://huggingface.co/datasets/ulamai/UnsolvedMath/blob/d371ee917ac9647fbb06cd6fc529deec6f3dda82/dataset.json.

[15] L. Bodnár, J. León, X. Liu and O. Pikhurko, *The Turán density of the tight 5-cycle minus one edge*, arXiv:2412.21011v3 (July 9, 2025; first posted December 30, 2024). https://arxiv.org/abs/2412.21011v3.

[16] J. Balogh and H. Luo, *Turán density of long tight cycle minus one hyperedge*, arXiv:2303.10530v2 (March 3, 2024; first posted March 19, 2023). https://arxiv.org/abs/2303.10530v2.

[17] B. Lidický, C. Mattes and F. Pfender, *The Hypergraph Turán Densities of Tight Cycles Minus an Edge*, arXiv:2409.14257v2 (November 26, 2024; first posted September 21, 2024). https://arxiv.org/abs/2409.14257v2.

[18] A. A. Razborov, *Flag algebras*, Journal of Symbolic Logic **72** (2007), no. 4, 1239–1282. https://doi.org/10.2178/jsl/1203350785.

[19] A. A. Razborov, *On 3-Hypergraphs with Forbidden 4-Vertex Configurations*, 2010. https://doi.org/10.1137/090747476. The earlier upper bound is cited through the text of [3, §2.4]; the original paper was not read in full.